



## BioFlora® Zinc (7%)

## Product Technical Data

Soil applications of **BioFlora® Zinc** are recommended with nitrogen fertilizer solutions. Thorough coverage and wetting are needed for optimum foliar spray results. Immediate post-irrigation or post rainfall applications give best foliar uptake. Addition of urea (for example, 5 lbs. Lo<Bi urea per 1000 gallons of spray solution) will enhance foliar uptake. Apply the recommended amount of product in split applications 1-2 weeks apart throughout the growing season. Contact a BioLynceus® representative for application intervals.

### Guaranteed Analysis

Zinc (Zn) 7%

### Derivation

Derived from zinc sulfate.

### Density

10.70 lb per gallon

1.28 kg per liter

### Application Details

Fertility Maintenance.....1-3 qts/acre

Mild Deficiency.....1-3 qts/acre

Severe Deficiency.....2-6 qts/acre

The lower rates are for field, row and vegetable crops. The higher rates are for fruit, tree, vine, ornamental and turf crops.

### Foliar Application

Apply 2-4 qts/acre as needed. Avoid foliar applications of this product when plants are under moisture stress. Avoid foliar applications to deciduous tree crops during bloom.

### Compatibility

BioFlora® Zinc is compatible with most pesticides and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past. Always add this product to spray tank before adding pesticides.



### **Product Support**

Consult your BioLynceus® representative for product support, application information and current pricing. Do not allow product to freeze.

### **Keep Away From Children – Not for Human Consumption**

*Note: buyer assumes all risk of use, storage and handling of this material. Neither the manufacturer, seller, nor its agents, make any warranty, expressed or implied, concerning this product, except in conformity with statements on the label. Information regarding this product can be obtained from BioLynceus®. Do not allow product to freeze.*